

LANDELEK



COMPOSITION

Nutrient	g/kg
Crude Protein (min.)	250
Equiv. Crude Protein from NPN (max.)	41.58 %
Urea (max.)	20
Moisture (max.)	120
Crude Fat (min.)	14
Crude Fibre (max.)	80
Calcium (min.)	8.1
Calcium (max.)	20
Phosphorus (min.)	6.5
Ca: P Ratio	1-3:1
Magnesium (min.)	2.5
Selenium (min.)	0.3 mg/kg
Copper (min.)	26 mg/kg
Iron (min.)	57 mg/kg
Manganese (min.)	40 mg/kg
Zinc (min.)	160 mg/kg
Cobalt (min.)	0.5 mg/kg
Iodine (min.)	1.3 mg/kg
ME Sheep	7.5 MJ/kg

FEEDING RECOMMENDATIONS

- This is a supplement and not a feed.
- Sufficient grazing must be available at all times.
- Keep lick troughs filled to prevent gluttonous eating by hungry animals.
- A constant daily intake can help prevent poisoning.
- A salt phosphorus lick must be available at least 7 days prior to feeding a NPN containing lick to prevent over consumption of urea due to salt hunger.
- Store in a cool, dry place, away from direct sunlight.

Form: Mash
Packaging: 40 kg Bags

RECOMMENDED INTAKE

Animal	g/animal/day
Sheep	300 - 400
Cattle	1200 - 1500

WARNING

This farm feed contains a NPN source and registered stock remedies. Feed strictly according to instructions.

12 Byls Bridge Boulevard
Highveld Ext 73
Centurion

Tel: +27 11 063 2347
afgrianimalfeeds.co.za

Ready-mix high quality production lick for cattle and sheep

- Suitable for sheep and cattle in production on low quality grazing or roughages.
- Provides optimal quantities of quality natural protein, NPN, minerals, trace minerals and energy for economic utilization of low quality roughages.
- Supplies high levels of rumen bypass protein and energy for improved ADG as well as milk and wool production.
- Recommended for flush feeding, pregnant and lactating animals and young stock.

FEEDING INSTRUCTIONS

1. This farm feed should not be fed indiscriminately with other NPN containing farm feeds. Consult an AFGRI Animal Feeds Technical Advisor for recommendations.
 2. Vinegar is an effective remedy against NPN poisoning. Treatment: Mix with an equal amount of water and dose half a bottle per calf or large sheep and 2 to 4 bottles per head of cattle (1 bottle = 750ml).
 3. Protect this farm feed against rain. NPN (e.g. urea) is soluble and animals drinking such a solution could be poisoned.
 4. Guidelines for intake: Intake will depend on the production status of the animals and the quality and quantity of pasture available.
 5. Lick intake needs to be managed. Ideally supply the required amount (g/animal/day) on a daily basis. It will not be a problem if the required amount is consumed within a few hours. If management practice only allows the supply of lick every third day, then the required amount of lick can be supplied (g/animal x number of animals x 2 days) every third day. Research shows it will not be a problem if animals consume the supplement in a shorter period. These feeding methods ensure accurate intake under all circumstances, which would not be possible if salt is used solely as an intake regulator.
- Sufficient trough space is of the utmost importance.**
6. An adaptation period of 14 days with maize is recommended, before permanent grazing of harvested crops, to prevent rumen acidosis for sheep.

ADAPTATION FEEDING GUIDELINES

An adaptation period of 14 days is recommended.

Day	Maize g/sheep/day	Grazing time/ day
Day 1-2	200	15 min
Day 3-4	300	30 min
Day 5-6	400	45 min
Day 7-8	500	60 min
Day 9-10	600	75 min
Day 11-12	700	90 min
Day 12-14	800	120 min

Class: PROTEIN, ENERGY, MINERAL AND TRACE
MINERAL SUPPLEMENT FOR SHEEP
Reg. No: V26177 – Act 36/1947
AFGRI Product code: R1216



ANIMAL FEEDS

LANDELEK



SAMESTELLING

Nutriënt	g/kg
Ruproteïen (min.)	250
Ekwivalente Ruproteïen vanaf NPN (maks.)	41.58 %
Ureum (maks.)	20
Vog (maks.)	120
Ru-vet (min.)	14
Ruvesel (maks.)	80
Kalsium (min.)	8.1
Kalsium (maks.)	20
Fosfor (min.)	6.5
Ca:P Verhouding	1-3:1
Magnesium (min.)	2.5
Selenium (min.)	0.3 mg/kg
Koper (min.)	26 mg/kg
Yster (min.)	57 mg/kg
Mangaan (min.)	40 mg/kg
Sink (min.)	160 mg/kg
Kobalt (min.)	0.5 mg/kg
Jodium (min.)	1.3 mg/kg
ME Skaap	7.5 MJ/kg

VOER AANBEVELINGS

- Die produk is 'n supplement en nie 'n voer nie.
- Voldoende weiding/ruvoer moet deurgaans beskikbaar wees.
- Hou lekbakke vol om gulsige vreet deur honger diere te voorkom.
- 'n Gereelde daaglikse inname kan vergiftiging help voorkom.
- 'n Soutfosfaatlek moet ten minste 7 dae voor die verskaffing van 'n NPN-bevattende lek voorsien word om southonger en oormatige inname van ureum te voorkom.
- Stoor in 'n koel, droë plek, weg van direkte sonlig.

Vorm: Meel
Verpakking: 40 kg Sakke

AANBEVOLE INNAME

Dier	g / dier / dag
Skaap	300 - 400
Beeste	1200 - 1500

WAARSKUWING

Hierdie veevoer bevat 'n NPN bron en geregistreerde medikamente.
Voer streng volgens instruksies.

12 Byls Bridge Boulevard
Highveld Ext 73
Centurion

Tel: +27 11 063 2347
afgrianimalfeeds.co.za

Klaar gemengde hoë kwaliteit produksielek vir beeste en skape

- Geskik vir produserende skape en beeste op lae kwaliteit weidings of ruvoer.
- Voorsien optimale kwantiteit en kwaliteit natuurlike proteïen, NPN, minerale, spoorminerale en energie vir ekonomiese benutting van lae kwaliteit ruvoere.
- Voorsien hoë vlakke van rumen verbyvloei proteïen en energie vir beter GDT asook melk- en wolproduksie.
- Word aanbeveel vir prikkelvoeding, dragtige- en lakterende diere en uitgroei van jong diere.

GEBRUIKSAANWYSINGS

1. Moet nie hierdie veevoer onoordeelkundig saam met ander NPN bevattende veevoere voer nie. Raadpleeg 'n AFGRI Veevoere Tegniese Adviseur vir die nodige aanbevelings.
2. Asyn is 'n doeltreffende middel teen NPN-vergiftiging. Behandeling: Meng met gelyke hoeveelhede water en doseer 'n halwe bottel per kalf of groot skaap en 2 tot 4 bottels per bees (1 bottel = 750ml).
3. Beskerm hierdie veevoer teen reën. NPN (Bv. Ureum) is oplosbaar en diere wat so 'n oplossing drink kan vergiftig word.
4. Riglyne vir inname: Inname sal bepaal word deur die produksie status van diere en die kwaliteit en kwantiteit van die weiding.
5. Lek inname moet bestuur word. Dit word aanbeveel om die verlangde hoeveelheid daaglik aan diere te voorsien. Navorsing toon egter dat indien diere die verlangde daaglikse hoeveelheid lek in 'n korter tyd vreet, daar nie probleme onder vind word nie. Indien bestuur slegs toelaat om bv. elke derde dag lek uit te plaas, verskaf dan die verlangde hoeveelheid lek as volg (g/dier x aantal diere x 2 dae). Hierdie voermetodes maak dit moontlik om onder alle omstandighede die verlangde innames te verkry, wat nie moontlik sou wees indien sout uitsluitlik as inname-reguleerder gebruik word nie. **Voldoende kripspasie is van uiterste belang.**
6. Skape moet vir 14 dae voordat hulle op mielielande gaan, op mielies aangepas word om suurpens te voorkom.

RIGLYNE VIR INNAME AANPASSING

'n Aanpassings periode van 14 dae word aanbeveel.

Dag	Mielies g/skaap/dag	Wei tyd/ dag
Dag 1-2	200	15 min
Dag 3-4	300	30 min
Dag 5-6	400	45 min
Dag 7-8	500	60 min
Dag 9-10	600	75 min
Dag 11-12	700	90 min
Dag 12-14	800	120 min

Klas: PROTEÏEN, ENERGIE, MINERAAL EN SPOORMINERAAL SUPPLEMENT VIR SKAPE
Reg. No: V26177 – Wet 36/1947
AFGRI Produktkode: R1216



ANIMAL FEEDS